

GEORGIA TECH

SCHOOL OF MATHEMATICS

MATH 1502D

CALCULUS II, SECTION D

Test # 1

September 22nd, 2010

First Name : -----

Last Name : -----

DO NOT WRITE IN THE TABLE BELOW

1a	
1b	
2a	
2b	
2c	
2d	
3	
4a	
4b	
4c	
4d	
5a	
5b	
6a	
6b	

WARNING :

Read carefully, read the comments in *italic*, take your time, do not panic and double check what you write.

Take the time to write in plain English the criteria or the names of the tests you are using to justify your answer.

The test will last 50 minutes.

1. (a) Give the Taylor *expansion* of $P(x) = 27 - 27x + 9x^2 - x^3$ near $x = 3$

$$P(x) =$$

- (b) Give the value of $Q^{(19)}(0)$ if

$$Q(x) = 1 - \frac{x^{13}}{39} + \frac{x^{19}}{17 \times 18 \times 19} - \frac{x^{39}}{37 \times 38 \times 39} + \frac{x^{52}}{437} - \frac{x^{65}}{65 \times 64 \times 63}$$

$$Q^{(19)}(0) =$$

2. The hyperbolic cosine and sine are defined by $\cosh x = \frac{e^x + e^{-x}}{2}$ and $\sinh x = \frac{e^x - e^{-x}}{2}$.

(a) Compute the first and the second derivative of $\cosh x$

$$\frac{d \cosh x}{dx} =$$

$$\frac{d^2 \cosh x}{dx^2} =$$

(b) Give the Taylor *expansion* to order $2n + 1$, near $x = 0$ of $\cosh x$

Hint : compute first $\cosh 0$ and $\sinh 0$

$$\cosh x =$$

- (c) Give the expression R_{2n+1} of the remainder (*Taylor form*) and show that $0 \leq R_{2n+1} \leq (\cosh x)x^{2n+2}/(2n+2)!$

(*Hint : remark that $\cosh x$ is positive, increasing in x for $x > 0$ and that $\cosh -x = \cosh x$*)

Put the result here and use the back page for your calculations

$$R_{2n+1}(x) =$$

- (d) Compute the numerical value of $\cosh 1$ by using $n = 2$: using the previous bound on the remainder gives an interval of values to which the exact value belongs.

(*Hint : $37/(24 \times 719) \leq 2.2 \times 10^{-3}$.)* **Put the result here and use the back page for your calculations**

$$\leq \cosh 1 \leq$$

(Use this page for your calculations)

3. Is the following integral convergent?

(Hint : identify the points on the interval of integration where the integral is improper; it will be admitted that $x^6 - x^3 + 1 \geq 3/4$)

$$\int_0^{\infty} \frac{dx}{(x^7 - x^4 + x)^{3/14}}$$

4. Tell whether the following series converge or not and indicate the tests used to conclude

(a)

$$\sum_{n=0}^{\infty} (-1)^n \left(\frac{n}{1+n} \right)^{31}$$

Converges

Diverges

Test used :

(b)

$$\sum_{n=0}^{\infty} \frac{167^n}{(n!)^{0.01}}$$

Converges

Diverges

Test used :

(c)

$$\sum_{n=0}^{\infty} \frac{\{\ln(n+2)\}^{153}}{(1+2n^3)^{1/2}}$$

Converges

Diverges

Tests used :

(d)

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{5n^2-1}}$$

Converges absolutely :

YES

NO

Converges

Diverges

Tests used :

5. Let $f(x)$ be the function given by the power series

$$f(x) = \sum_{n=0}^{\infty} \frac{x^{2n}}{(1 + 5n + n^2)}$$

- (a) Compute the radius of convergence and indicate the test you are using.

Domain of convergence =

Arguments and Tests used :

- (b) What may happen at the end points?

End points :

(Use this page for your calculations)

6. This question concerns the convergence of the following series

$$\sum_{n=1}^{\infty} \frac{n^2}{1 + 2^3 + 3^3 + \cdots + n^3} \quad (1)$$

- (a) Using a comparison with the function $f(x) = x^3$, show that $1 + 2^3 + 3^3 + \cdots + n^3 \geq \int_0^n x^3 dx$

- (b) Using the previous estimate, is the series (1) above converging or diverging? Give the criterion used in to justify your answer.

(Use this page for your calculations)